

Tetanus after Trauma: Still Existed Problem in Tropical Country

Viroj Wiwanitkit

Department of Tropical Medicine, Hainan Medical University, Haikou, China

Address for correspondence: Prof. Viroj Wiwanitkit,
Hainan Medical University, Haikou, China.
E-mail: wviroj@yahoo.com

Dear Editor,

An important concern on the management of trauma case is the tetanus prevention. In the past, when tetanus toxoid, is not easily available, the rate of posttraumatic tetanus was very high. In tropical countries, the problem of tetanus is still existed. Here, the author discusses and summarizes the problem of tetanus after trauma in the local reports^[1,2] in Thailand, a tropical country in Asia. Based on database review (PubMed, Scopus, and Thai Index Medicus) on case reports of tetanus in Thailand in the past 15 years period (2001–2016), there are at least 132 accumulated cases of posttraumatic tetanus. The male-to-female ratio was 2:1. The mortality rate is 8.02%. In all cases, the patients did not receive the tetanus toxoid.

It is interesting that tetanus still exists in the period after the millennium, the year 2000 in a tropical country where the sanitation and public health system is significantly better at present. The history tracking shows that almost all patients neglected the importance of posttraumatic tetanus vaccination due to the small wound. Nevertheless, there are also some cases of minorities who cannot assess the medical facilities to get vaccinated. The importance to promote the knowledge of wound care and posttraumatic tetanus vaccination to the local people is required. As noted by Pandejpong and Phisalprapa,^[3] “*vaccination program remains a problem in Thailand especially among older generations and low-socioeconomic groups.*” Furthermore, the coverage of service to the minorities who lack opportunities to assess medical facilities is needed.^[4]

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Sathirapanya P, Sathirapanya C, Limapichat K, Setthawacharawanich S, Phabphal K. Tetanus: A retrospective study of clinical presentations and outcomes in a medical teaching hospital. *J Med Assoc Thai* 2009;92:315-9.
2. Kanchanapongkul J. Tetanus in adults: A review of 85 cases at Chon Buri Hospital. *J Med Assoc Thai* 2001;84:494-9.
3. Pandejpong D, Phisalprapa P. Awareness of and factors influencing willingness to receive a tetanus-diphtheria toxoid booster vaccine among Thai adults. *Southeast Asian J Trop Med Public Health* 2013;44:1047-54.
4. Greenfield LS, Page LC, Kay M, Li-Vollmer M, Breuner CC, Duchin JS, *et al.* Strategies for increasing adolescent immunizations in diverse ethnic communities. *J Adolesc Health* 2015;56:S47-53.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:



Website:

www.archtrauma.com

DOI:

10.4103/atr.atr_59_17

How to cite this article: Wiwanitkit V. Tetanus after Trauma: Still Existed Problem in Tropical Country. *Arch Trauma Res* 2017;6:105.

© 2018 Archives of Trauma Research | Published by Wolters Kluwer - Medknow